

The following table shows the results of the regression analysis for the dependent variable *Y* (in thousands of dollars) against the independent variable *X* (in thousands of dollars). The regression equation is  $\hat{Y} = 1.2X + 0.5$ . The coefficient of determination is  $R^2 = 0.85$ .

✓

1. A method for treating cancer in a mammal, said method comprising:  
administering to a tumor site of the mammal an anticancer composition  
comprising a mixture of an anticancer agent and a calcium phosphate paste, said  
paste comprised of one or more nanocrystalline or poorly crystalline calcium  
5 phosphates and a physiologically acceptable fluid, the paste having an injectable or  
formable consistency at the time of administration and hardenable at the tumor  
site.

2. The method of claim 1, wherein each calcium phosphate having a  
Ca/P ratio of less than or equal to 1.7.

3. The method of claim 1, wherein the anticancer agent is selected from  
the group consisting of methotrexate, cis-platin, prednisone, hydroxyprogesterone,  
medroxyprogesterone acetate, megestrol acetate, diethylstilbestrol, testosterone  
propionate, fluoxymesterone, vinblastine, vincristine, vindesine, daunorubicin,  
doxorubicin, hydroxyurea, procarbazine, aminoglutethimide, mechlorethamine,  
15 cyclophosphamide, mephalan, uracil mustard, chlorambucil, busulfan, carmustine,  
lomusline, dacarbazine (DTIC, dimethyltriazenomideazolecarboxamide),  
fluorouracil, 5-fluorouracil, cytarabine, cytosine arabinoside, mercaptopurine, 6-  
mercaptopurine, tamoxifen, paclitaxel, etoposide, vinorelbine, gemcitabine,  
leuprolide, flutamide, goserelin acetate, and thioguanine, and mixtures thereof.

4. The method of claim 1, wherein the anticancer composition is  
administered to the tumor site by cannula or by injection.

5. The method of claim 4, wherein the anticancer composition is administrable by cannula or injection more than five minutes after its preparation.

6. The method of claim 5, wherein the anticancer composition is administrable by cannula or injection more than twenty minutes after its preparation.

7. The method of claim 1, wherein the paste hardens into an apatitic calcium phosphate.

8. The method of claim 1, wherein the nanocrystalline or poorly crystalline calcium phosphate paste comprises a calcium phosphate selected from the group consisting of poorly crystalline apatitic (PCA) calcium phosphates (PCA), dicalcium phosphates, such as dicalcium phosphate dihydrate (DCPD) and dicalcium phosphate anhydrous (DCPA), tricalcium phosphates (TCP), monetite, monocalcium phosphate monohydrate (MCPM), hetpacalcium phosphate, calcium pyrophosphate, calcium metaphosphate, octacalcium phosphates (OCP), hydroxyapatites (HA).

9. The method of claim 8, wherein at least one of the nanocrystalline or poorly crystalline calcium phosphates is a poorly crystalline apatitic calcium phosphate.

10. The method of claim 1, wherein each of the said one or more nanocrystalline or poorly crystalline calcium phosphates has a calcium to

phosphate ratio in the range of 1.0 to 1.67.

11. The method of claim 1 wherein each of the said one or more nanocrystalline or poorly crystalline calcium phosphates has a calcium to phosphate ratio in the range of 1.3 to 1.67.

5 12. The method of claim 1 wherein the nanocrystalline or poorly crystalline calcium phosphate paste has an overall calcium to phosphate ratio in the range of 1.0 to 1.7.

10 13. The method of claim 1, wherein the nanocrystalline or poorly crystalline calcium phosphate paste has an overall calcium to phosphate ratio in the range of 1.40 to 1.65.

14. The method of claim 1, wherein the nanocrystalline or poorly crystalline calcium phosphate paste comprises a physiologically acceptable fluid in an amount sufficient to produce a paste having injectable or formable consistency.

15 15. The method of claim 1, wherein a therapeutically effect amount of anticancer agent is released from the composition for a time greater than one week.

16. The method of claim 1, wherein a therapeutically effect amount of anticancer agent is released from the composition for a time greater than two week.

17. The method of claim 1, wherein a therapeutically effect amount of

anticancer agent is released from the composition for a time greater than one month.

18. The method of claim 1, wherein a therapeutically effect amount of anticancer agent is released from the composition for a time greater than three  
5 months.

19. The method of claim 1, wherein delivery of the anticancer therapy to the tumor site is sufficient to prevent increase of tumor mass without significant weight loss of the mammal.

10 20. The method of claim 1, wherein delivery of the anticancer therapy to the tumor site is sufficient to result in a decrease in tumor mass without significant weight loss in the mammal.

15 21. The method of claim 1, wherein the particle size of the nanocrystalline or poorly crystalline calcium phosphate is selected to provide a desired release kinetic of the anticancer drug.

Sub B1  
20 22. An anticancer composition comprising a mixture of a physiologically effective amount of an anticancer agent and a calcium phosphate paste, said paste comprised of one or more nanocrystalline or poorly crystalline calcium phosphate and a physiologically acceptable fluid, the paste having an injectable or formable consistency at the time of administration and hardenable at the tumor site.

23. The composition of claim 22, wherein each calcium phosphate having a Ca/P ratio of less than or equal to 1.7.

SUB  
A1

24. The composition of claim 22, wherein the anticancer agent is selected from the group consisting of methotrexate, cis-platin, prednisone, hydroxyprogesterone, medroxyprogesterone acetate, megestrol acetate, diethylstilbestrol, testosterone propionate, fluoxymesterone, vinblastine, vincristine, vindesine, daunorubicin, doxorubicin, hydroxyurea, procarbazine, aminoglutethimide, mechlorethamine, cyclophosphamide, mephalan, uracil mustard, chlorambucil, busulfan, carmustine, lomusline, dacarbazine (DTIC, dimethyltriazenomideazopolecarboxamide), fluorouracil, 5-fluorouracil, cytarabine, cytosine arabinoside, mercaptopurine, 6-mercaptopurine, tamoxifen, paclitaxel, etoposide, vinorelbine, gemcitabine, leuprolide, flutamide, goserelin acetate, and thioguanine, and mixtures thereof.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

25. The composition of claim 22, wherein the anticancer composition is of a consistency administrable to the tumor site by cannula or by injection.

SUB  
A1

26. The composition of claim 22, wherein the nanocrystalline or poorly crystalline calcium phosphate cement comprises a calcium phosphate selected from the group consisting of amorphous calcium phosphate, poorly crystalline apatitic (PCA) calcium phosphates (PCA), dicalcium phosphates, such as dicalcium phosphate dihydrate (DCPD) and dicalcium phosphate anhydrous (DCPA), tricalcium phosphates (TCP), monetite, monocalcium phosphate monohydrate (MCPM), heptacalcium phosphate, calcium pyrophosphate, calcium

SUB  
A2

metaphosphate, octacalcium phosphates (OCP), hydroxyapatites (HA).

27. The composition of claim 26, wherein at least one of the nanocrystalline or poorly crystalline calcium phosphates is selected from the group consisting of amorphous calcium phosphate and poorly crystalline apatitic calcium phosphate.

28. The composition of claim 22, wherein each of the said one or more nanocrystalline or poorly crystalline calcium phosphates has a calcium to phosphate ratio in the range of 1.3 to 1.67.

29. The composition of claim 22, wherein the nanocrystalline or poorly crystalline calcium phosphate paste has an overall calcium to phosphate ratio in the range of 1.0 to 1.7.

30. The composition of claim 22, wherein the nanocrystalline or poorly crystalline calcium phosphate paste has an overall calcium to phosphate ratio in the range of 1.0 to 1.67.

31. The composition of claim 22, wherein the nanocrystalline or poorly crystalline calcium phosphate paste has an overall calcium to phosphate ratio in the range of 1.40 to 1.65.

32. The composition of claim 22, wherein nanocrystalline or poorly crystalline calcium phosphate paste comprises a physiologically acceptable fluid in

an amount sufficient to produce a paste having injectable or formable consistency for at least five minutes.

33. The composition of claim 22, wherein nanocrystalline or poorly crystalline calcium phosphate paste comprises a physiologically acceptable fluid in  
5 an amount sufficient to produce a paste having injectable or formable consistency for at least twenty minutes.

34. The composition of claim 22, wherein the nanocrystalline or poorly crystalline calcium phosphate paste is hardenable into an apatitic calcium phosphate.

35. The composition of claim 22, wherein a therapeutically effect amount of anticancer agent is released from the composition for a time greater than one week.

SUB  
A3

36. The composition of claim 22, wherein a therapeutically effect  
15 amount of anticancer agent is released from the composition for a time greater than two week.

37. The composition of claim 22, wherein a therapeutically effect amount of anticancer agent is released from the composition for a time greater than one month.

20 38. The composition of claim 22, wherein a therapeutically effect



amount of anticancer agent is released from the composition for a time greater than three months.

39. The composition of claim 22, wherein delivery of the anticancer  
5 therapy to the tumor site is sufficient to at least prevent increase of tumor mass without significant weight loss of the mammal.

*Sub 3*  
40. The composition of claim 22, wherein delivery of the anticancer  
therapy to the tumor site is sufficient to prevent a decrease in tumor mass without  
significant weight loss in the mammal.

*Sub 3*  
41. The composition of claim 22, wherein the particle size of the  
10 calcium phosphate is selected to provide a desired release kinetic of the anticancer drug.

*Sub 4*  
42. A kit for use in preparing a flowable anticancer composition that  
remain injectable for at least about 20 minutes, said kit comprising:

dry ingredients comprising a nanocrystalline or poorly crystalline  
calcium phosphate and a second calcium phosphate in a proportion of about 1:10  
to 10:1 by weight;

a physiologically acceptable aqueous lubricant in an amount sufficient  
to produce a flowable product upon combination with said dry ingredients; and

20 an anticancer agent in an amount ranging from about 0.01 to 10 wt. % of  
said dry ingredients.

